

Treatment of Prostate Cancer with Natural Therapeutics

— THIRD EDITION —
SEPTEMBER 2002

Published as a public service by **Prostate Cancer Fund**
(*A special program of Project Cure*)
P.O. Box 96673, Washington, D.C. 20090-6673 • 1-800-716-2152

Introduction:

The prostate is a gland in the male reproductive system that helps produce semen, the thick fluid that carries sperm cells. The prostate is a walnut-sized structure located beneath the bladder of males. It surrounds the upper part of the urethra. The urethra is the tube that carries urine from the bladder. Prostate function is regulated by testosterone, the male sex hormone produced primarily in the testicles.

Prostate cancer represents a significant number of malignancies in men over the age of 50 years. The incidence of prostate cancer has been increasing each decade in the United States. About 200,000 new cases and 38,000 deaths are related to prostate cancer each year in the United States. African-American males have the highest incidence of prostate cancer in the world.¹ The disease is rare in Asia, Africa, and Latin America.

Prostate cancer typically has a strange course. It often is inactive for many years. Once activated, it tends to progress slowly. The disease is often fatal if it spreads beyond the prostate gland. The cause of prostate cancer is still unknown, but it is likely related to hormonal factors. In particular, the conversion of testosterone to dihydrotestosterone (DHT) is recognized as an important factor in the development of disease. Adenocarcinoma is the primary type of prostate cancer, but sarcoma, squamous cell carcinoma, ductal transitional carcinoma, and undifferentiated prostatic carcinoma also occur.

Signs and Symptoms:

Prostate cancer is typically a progressive disease that occurs at a slow rate. It can also develop with no significant symptoms, particularly in the early stages of the disease. As the disease progresses, however, symptoms such as the following begin to appear:

- The need to urinate frequently, particularly at night
- Problems in the initiation and cessation of urination
- The complete inability to urinate
- Blood in the urine
- Painful ejaculation
- Burning or painful urination
- Persistent pain in the upper thighs, lower back, or hips

It is important to note that benign conditions involving the prostate such as enlargement (benign prostatic hypertrophy, BPH), infections, and stones can mimic the symptoms of cancer, so a professional

consultation is important. When the cancer metastasizes to the ribs, pelvis, or vertebral bodies it can lead to bone pain in prostate cancer patients.

Diagnosis

Prostate cancer diagnosis begins with a digital rectal examination by the health care professional. Additional tests such as a transrectal or transperineal needle biopsy can detect small prostatic nodules or more advanced cancerous growth. Transrectal ultrasound is also used for both diagnosis and to help guide needle biopsy. Serum acid phosphatase levels can be elevated when there is local extension or metastasis of disease. However, other conditions such as benign prostatic hyperplasia, multiple myeloma, Gaucher's disease, and hemolytic anemia can also raise these levels. Radioimmunoassay analysis of prostate-specific antigen (PSA) has become a standard test for diagnosing both localized as well as metastasized stages of the prostate cancer. Though controversial, PSA is the most sensitive marker for monitoring disease and is elevated in 25 to 92% of patients with prostate cancer. However, it is also raised in 30 to 50% of those with BPH. Prostate cancer often leads to osteoblastic (bone-formation cells) bony metastases, which are often detected on bone scans or x-rays of prostate cancer patients. Physicians often have difficulty in identifying what types of prostate cancers will spread and become dangerous. A new protein called KAI-1 has been identified as a marker for disease. If significant levels of KAI-1 are found in cancerous tissues, the cancer is not likely to spread. If the protein is not found, cancer typically spreads.

Prostate cancer has been classified into four stages:

- Stage A. The tumor is not found by normal tests but during surgery for a different prostate condition.
- Stage B. The tumor can be palpated during rectal examination but it has not spread beyond the prostate
- Stage C. Cancer has spread to tissues outside of the prostate.
- Stage D. The cancer has metastasized to the lymph nodes of the pelvis or to other parts of the body, especially the bones.

Conventional Therapy

Prostate cancer that is localized is typically treated with radiation therapy or radical prostatectomy (excision of part or all of the prostate gland). This operation is performed through incision in the perineum, into the bladder, or through the urethra.

Libido is typically unaffected by prostatectomy but some 30% of patients become impotent after the procedure. A more common complication following a radical prostatectomy is the development of urinary incontinence. When disease has significantly progressed beyond this stage, or when the patient is very old or in poor health, these treatment approaches may not be used. These patients may be treated with irradiation, hormone therapy, or the surgical removal of the testicles. One of the hormonal therapies, oral diethylstilbestrol, has been given to prostate cancer patients in doses of 1 to 3 mg/day and has demonstrated some success over long periods of time. Long-term use of such estrogen therapies increases the risk of developing thromboembolic (blocking of a blood vessel by a thrombus) complications. Additional adverse effects caused by estrogen therapies can include breast tenderness, breast enlargement, nausea, vomiting, loss of sexual desire, impotence, and water retention. Short-term use of agents such as high-dose diethylstilbestrol diphosphate can lead to substantial relief in patients within days. A variety of agents are used to decrease testosterone levels circulating in the body. These agents include flutamide, cyproterone acetate, ketoconazole, aminoglutethimide, and analog agents of luteinizing hormone-releasing hormone. Surgical removal of the testicles is sometimes performed when the disease has advanced or when hormone therapy has failed. The use of local radiation therapy has been found to be effective in relieving pain associated with cancer metastasis into the bones. Local radiation therapy can also help limit disease to the prostate. Chemotherapy has generally not been effective once hormonal therapy has failed. It is also associated with severe adverse effects such as nausea, vomiting, lowered blood and immune system factors, and hair loss.

Benign Prostatic Hypertrophy

While BPH and prostate are not the same condition and are not believed to be related to each other, they do have many common factors. As mentioned previously, both produce elevated PSA levels and a variety of problems relating to urination. In addition, both are believed to develop from hormonally-related factors. BPH occurs in more than half of all men in their 60s, and in as many as 90 percent of all men in their 70s and 80s. BPH may develop from the growth of cells from the relatively increased levels of estrogen that occur in men as they age. Another line of thought states that the testosterone-derivative DHT⁹⁵ is involved with the increased cellular growth associated with BPH. The drug called finasteride (Proscar), a 5-alpha-reductase inhibitor has reduced the size of the prostate

in some patients, resulting in improved urination. BPH does not increase the risk of developing prostate cancer, and these two conditions develop in different parts of the prostate gland.

Risk Factors

Age, as previously mentioned, is a significant risk factor for disease development. Men with a family history of prostate cancer have a significantly increased risk for the development of the disease. A man whose father had prostate cancer is 2.3 times more likely to develop the disease than if his father never developed the disease. A man is also more likely to develop prostate cancer if his sister or mother develops breast or ovarian cancer. This suggests an important genetic relationship in the development of prostate cancer. Some 15% of inherited cases may be transmitted through the mother's X chromosome. A different gene called HPC1 has been associated with about 30% of all inherited cases of prostate cancer. Some association between viral and bacterial infections of the prostate and the later development of prostate cancer has been seen.

Levels of sexual activity do not appear to have a significant effect on the risk of developing disease. Despite some discussion to the contrary, there is no evidence that a vasectomy will alter prostate cancer risk. Occupational factors may be a small factor in the development of disease. There is some evidence that working around metals and chemicals, such as acrylonitrile, dimethylformamide, and cadmium, may increase the risk of developing prostate cancer. There is also limited evidence that a history of nonmelanoma skin cancer may increase the risk of developing disease.

Apart from genetic and age factors, the strongest evidence of risk association comes from dietary patterns. Research has suggested that men who consume high levels of fat are more likely to develop prostate cancer. Prostate cancer is more common in countries where meat and dairy products are consumed in large quantities than in regions where the diet contains large quantities of vegetables, rice, and soybean products. Fat has been found to stimulate testosterone and other hormone production. A recent study in Sweden found that a compound called acrylamide, which is found in many high-carbohydrate foods such as potato chips, fresh potatoes, and bread, is a powerful carcinogen.¹⁴⁹ Acrylamide, a substance used in water treatment facilities throughout the world, is apparently released in higher quantities when cooking or boiling carbohydrates. High testosterone

levels may also stimulate dormant prostate cancer cells into action. Eating meat cooked at high temperatures can produce carcinogens that directly affect the prostate. A recent study found that a gene that is necessary for the digestion of red meat is found in higher quantities in the prostatic tissue of patients with prostate cancer than in those with healthy prostate tissue.¹⁵⁰ Smokers who develop prostate cancer tend to have poorer outcomes than those who do not smoke.² Studies have also shown that the ingestion of 22 to 56 alcoholic drinks per week also increases risk of developing the disease.³

Obesity as a Risk Factor

Obesity has been associated with an increased risk of developing prostate cancer. A study of 5,212 men who died from prostate cancer and others who did not develop the disease found that obese (body mass index greater than 30) were 1.27 times more likely to develop and die from the disease than non-obese men.¹²⁸ A study from China found that men with the highest waist-to-hip ratios were nearly three times more likely to develop prostate cancer as those with the smallest waist-to-hip ratios.¹²⁹ This may be due to the fact that greater fat deposits in the abdomen may increase hormonal activity. A study of 101 men who developed prostate cancer in Iowa found that obesity was associated with more aggressive and late-stage prostate tumors.¹³⁰ A different study found an association between childhood obesity and an increased risk of prostate cancer late in life.¹³¹

Prostate Cancer Prevention

The prevention of prostate cancer has just begun to be studied. At the microscopic level, men throughout the world appear to have the early stage of the disease. However, the large variations in the progression of the disease among populations in particular regions is the focus of researchers. For example, Asians who move to the United States have a higher incidence of prostate cancer than those who live in Asia.⁴ This suggests a role for nutrition in the prevention of the disease. Animal fats and certain vegetable oils appear to be critical factors in disease development. Diets that involve the ingestion of large amounts of fresh fruits and dark-colored vegetables have been shown, in general, to have cancer-prevention effects.

Phytonutrients

Even though there is evidence that supplementation of various nutrients to the diet can provide benefits in the prevention and therapy of prostate cancer, more benefit may be derived from obtaining various nutrients

from natural sources in the diet. In fact, the National Cancer Institute recommends the ingestion of at least five servings daily of fruits and vegetables to fight the development and progression of prostate cancer.¹⁴⁵ In addition, scientists have discovered that not all nutrients have been identified in some foods. The phytochemicals known as silymarin, genistein, and epigallocatechin 3-gallate have been shown to inhibit prostate cancer cells from multiplying.¹⁴⁶ Several studies have found that vegetables, fruits, and whole grains contain numerous phytonutrients that modulate cancer development. These foods contain anticarcinogenic compounds such as chlorophyll, carotenoids, flavonoids, indole, polyphenols, and protease inhibitors.¹⁴⁷ Researchers have found that the foods and herbs with the highest anticarcinogenic activity are licorice, garlic, soybeans, cabbage, ginger, citrus, and the umbelliferous vegetables.¹⁴⁸

Lycopene

Lycopene is a chemical in some fruits and vegetables that has demonstrated some potential in the prevention of prostate cancer.⁵ Lycopene is being thoroughly studied in a number of ongoing trials. Lycopene is a carotenoid present in blood that has proven antioxidant activity.⁶ In vitro and in vivo studies have shown that lycopene has protective effects against some types of cancers. Lycopene ingestion has been shown to reduce some types of digestive system cancers but has been primarily studied in association with prostate cancer.⁷ Lycopene is found primarily in nature in tomatoes and tomato products but is also found in carrots, green peppers, apricots, watermelon, and pink grapefruit. A case-control study that analyzed plasma levels of lycopene in men with prostate cancer and in healthy men found that men with the highest levels of lycopene in plasma were less likely to develop prostate cancer.⁸ An additional study of 12 prostate cancer patients and 12 age-matched healthy subjects found significantly lower lycopene serum and tissue levels in cancer patients than in controls.⁹ A study of mortality from prostate cancer in 41 countries found that men who ingest more than 6 Kcal per day of tomatoes and tomato products have a significantly reduced risk of developing prostate cancer.⁵ A population-based case-control study of 317 prostate cancer cases and 480 controls in New Zealand found that the intake of lycopene and tomato-based foods was weakly associated with prostate cancer risk reduction.¹⁰ A prospective cohort study of 47,894 subjects found lycopene intake from tomato products was inversely associated with the risk of prostate cancer.¹¹ A study found that lycopene

obtained by eating tomato sauce is associated with a decreased risk of prostate cancer development.⁹⁵ A new study in mice found that lycopene has the ability to inhibit tumor formation in the prostate and that these effects are organ-specific.⁹⁶ It is generally believed the antioxidant effects of lycopene are responsible for the tumor-inhibiting properties of lycopene, but other mechanisms may also be involved.⁹⁷

A more recent study found that short-term supplementation with tomato sauce containing a particular form of lycopene (all-trans-lycopene) produces significant increases in total lycopene in serum and in the prostate gland.¹⁵¹ A recent review of the scientific literature found evidence that lycopene decreases the risk of developing prostate cancer.¹⁵² A prospective study recently found that two or more servings of tomato products or lycopene per week reduced the risk of developing prostate cancer by at least 23% compared to those who ate tomato products/lycopene less than once per month.¹⁵³ A study of 32 men with a history of prostate cancer and surgical prostate removal found that a diet that contains 30 milligrams of lycopene per day reduced oxidative DNA damage in prostate tissue by an average of 28 percent.¹⁵⁴ This regimen also reduced PSA levels by an average of 17.5 percent. This amount of lycopene is equivalent to about three-quarters of one cup of tomato sauce per day. A study from the University of California at Los Angeles found that a combination of lutein and lycopene reduced prostate cancer cell growth by 20 percent.¹⁵⁵ The commonly seen autumn olive shrub produces berries that contain 18 times more lycopene than tomatoes.

Selenium

Selenium has been shown to have anti-carcinogenic effects both in vitro and in clinical studies.¹² An in vitro study of both human and mouse tumor cell lines found that selenium reduced human prostate carcinoma cell growth by 50% compared to controls.¹³ A case-control study of 164 prostate cancer patients and 152 controls with BPH found that plasma selenium levels were significantly higher in the BPH group compared to the cancer group.¹⁴ High levels of selenium have been found to counteract the prostate cancer cell-stimulatory effects of cadmium in vitro.¹⁵ A case-control study of 181 prostate cancer cases and 181 matched controls found that selenium levels in the body, as gauged by levels in the toenails, are associated with the risk of prostate cancer.¹⁶ In these cases, high selenium levels were associated with significantly reduced risk of advanced prostatic cancer.

This held up even after controlling for a variety of factors, such as family history of prostate cancer, saturated fat intake, vasectomy, geographical region, body mass intake, and calcium intake. A study of 9345 Japanese-American men who had serum samples drawn and frozen in the 1970s and were assessed for prostate cancer incidence 20 years later found that those with selenium levels in the highest quartile were only half as likely to have disease as those in the lowest quartile.¹⁷ A similar study collected samples from 111 subjects in 1973 and found that the mean serum selenium levels of those with prostate cancer were significantly lower than those in 210 matched controls.¹⁸ A different study evaluated the effects of oral selenium supplementation on general cancer risk reduction in patients with a history of skin cancer.¹⁹ The researchers found no evidence that selenium supplementation affected skin cancer outcomes. However, they did find that supplementation reduced the incidence of all cancers combined, including lung, colorectal, and prostate. Selenium supplementation also reduced mortality from all cancers combined and total mortality. An additional study of 974 men with a history of basal or squamous cell carcinoma evaluated whether supplemental selenium affects the incidence of prostate cancer.²⁰ The men were randomized either to 200 micrograms of selenium daily or placebo for a mean length of 4.5 years. After an overall follow-up period of 6.5 years, the researchers found that the group receiving selenium had a significantly lower (63%) incidence of prostate cancer. The selenium group also had significantly lower total cancer incidence and overall mortality. A new study has shed some light on the mechanism by which selenium may produce some of its anti-tumor effects.⁹⁸ This study strongly suggests selenium plays a key role in the induction of apoptosis, or programmed cell death, of prostate cells. Several ongoing trials are evaluating the ability of selenium supplementation to reduce the risk of prostate cancer. A more recent study found that men with low serum levels of selenium have some reduction in prostate cancer risk if they use selenium supplementation.¹⁵⁶ A study in rats found that a diet containing broccoli sprouts and florets enriched with selenium reduced the risk of prostate cancer in treated animals.¹⁶³

Vitamin E

A study based on a cohort of 29,133 men found 317 developed prostate cancer during 9 years of follow-up. Baseline serum vitamin E levels were not important in prostate cancer development, but those who received high levels of vitamin E supplementation had a significantly reduced risk of developing prostate

cancer.²¹ Alpha-tocopherol combined with lycopene has been shown to have strong inhibitory effects on prostate cell proliferation in vitro.²² A new study has discovered that long-term vitamin E supplementation is associated with decreased levels of male hormones in the blood.⁹⁹ The lower level of hormone in the blood may be the critical factor in the found association between vitamin E supplementation and reduced risk of developing prostate cancer. A recently completed study of 145 men who developed prostate cancer found that these men had lower plasma levels of selenium, alpha-tocopherol (a component of vitamin E), and gamma-tocopherol (another component of vitamin E) than a control group that did not have the disease.¹⁰⁰ A new study discovered that vitamin E produces these anti-tumor effects on prostate cells by way of apoptosis (programmed cell death).¹⁰¹

A more recent study found that a form of vitamin E called vitamin E succinate kills prostate cancer cells but leaves normal prostate cell intact.¹⁵⁷ The recent Alpha-Tocopherol Beta-Carotene Prevention Study found that long-term supplementation with vitamin E reduces the risk of developing prostate cancer. A recent analysis of that study found that vitamin E lowered the level of male sex hormones, a key factor in the development of prostate cancer, in the serum of the treated group.¹⁵⁸ Vitamin E has been found to be an efficient regulator in the destruction of tumor cells that damage the ability of the body to fight cancerous growth.¹⁵⁹ Vitamin E appears to have especially strong anti-prostate cancer effects in younger men.¹⁶⁰ Vitamin E has also shown promise in the treatment of hot flashes in men who have undergone hormonal treatment for prostate cancer.¹⁶¹ A study has found that vitamin E interferes with the ability of certain cells to form receptors that are crucial in the development of prostate cancer.¹⁶²

Vitamin D

A study in rats found that an analog of vitamin D suppressed tumor development in the animals.²³ A different study in rats found that the administration of hormonal analog of vitamin D, 1, 25-dihydroxyvitamin D (calcitriol) and a different vitamin D analog called EB1089 led to significantly smaller prostate tumor volumes than a control solution.²⁴ An additional study also found evidence that 1,25-dihydroxyvitamin D inhibits the growth of both primary cultures of human prostate cancer cells and cancer cell lines.²⁵ Yet another study found that 1 alpha, 25-dihydroxyvitamin D can inhibit the invasive growth properties of the human prostate carcinoma lines, DU 145, PC-3, and LNCaP,

in vitro.²⁶ Epidemiological studies in humans have suggested vitamin D deficiency is an important factor in the development of prostate cancer.²⁷ A new study has found that prostate cancer cells lack a component of vitamin D that is typically present in healthy prostate cells.¹⁰² The same component found in this study, 1 alpha,25(OH)(2)D(3), may be a useful prostate cancer treatment when combined with previously-used therapies.¹⁰³ Another new study has found further evidence that this component of vitamin D can inhibit the growth of prostate cancer cells.¹⁰⁴

Recent research suggests a compound related to vitamin D called 1 alpha, 25 Dihydroxyvitamin D(3) (calcitriol) can alter certain mechanisms within cells in the prostate and produce increased cancer-fighting effects.¹⁶⁴ Calcitriol is the active metabolite of vitamin D in the body. Other research has found that calcitriol can maintain vitamin D levels in men being treated for prostate cancer.¹⁶⁵ These men often have significant vitamin D depletion during therapy, and this reduces the effectiveness of treatment. A recent study of 11 men being treated for prostate cancer found that combination treatment of docetaxel and calcitriol led to significant improvements in disease in all five men who completed eight weeks of therapy.¹⁶⁶ Docetaxel is a commonly used chemotherapeutic drug. A different but related compound of vitamin D called analogue (V) has been found to decrease proliferation of a prostate cancer cell line called DU145.¹⁶⁷ A recent review of studies involving the use of calcitriol on prostate cancer cells found significant evidence of anti-cancer effects.¹⁶⁸ Researchers previously believed that calcitriol required male sex hormones to produce its anti-cancer effects, but a recent study found that this is not true.¹⁶⁹ This may be an important consideration for treatment because many men with prostate cancer are treated with male hormone suppression therapy. The combination of calcitriol with a viral therapy produced powerful anti-prostate cancer cell effects in a recent study.¹⁷⁰

Zinc

Zinc plays a major role in androgen metabolism. Estrogens appear to reduce the uptake of zinc in the intestines. Estrogen levels are increased in men with BPH and sometimes in men with prostate cancer. Zinc has been shown to decrease the size of the prostate and to decrease symptomology in BPH patients.^{28,29} This could have implications for prostate cancer prevention and the patient recovery after prostatectomy. Zinc has also been shown to inhibit 5-alpha-reductase activity, which reduces the conversion of testosterone

to the dangerous DHT form.^{30,31} Zinc, like saw palmetto, has been shown to prevent the specific binding of androgens to nuclear and cytosol androgen receptors.³² Zinc has also been shown to inhibit prolactin secretion.³³ As mentioned before, prolactin increases the uptake of testosterone by the prostate. This increased prolactin secretion leads to greater levels of DHT because of increased levels of substrate.³⁴ Cadmium is an antagonist of zinc and is also a stimulant of 5-alpha-reductase activity. Zinc has been shown to selectively inhibit prolactin secretion from the pituitary.³⁵ Prolactin secretion directly stimulates increased production of testosterone and the conversion to the more dangerous form, DHT. A new study has determined that zinc is absolutely vital to the apoptosis of prostate cancer cells through its effects on the compound known as fetuin.¹⁰⁵ A different study discovered that zinc plays an important role in the regulation of tumor cell invasion of various tissues in the prostate.¹⁰⁶ An additional study confirmed these tumor-inhibiting effects of zinc.¹⁰⁷

A recent study found that zinc is involved in the regulation of enzymes that are involved with the invasion and metastasis of prostate cancer cells.¹⁷¹ A study found that zinc plays a vital role in the ability of a substance called fetuin to induce apoptosis, or cell death, in a variety of cancer cell types.¹⁷² Fetuin itself has been found to inhibit prostate cancer development in mice. A zinc-controlled factor called Kruppel-like factor 6 (KLF6) is mutated in a significant number of prostate cancer cases.¹⁷³ Zinc has been found in decreased levels in the cytoplasm of prostate cancer cells when compared to levels in the cytoplasm of normal prostate cells.¹⁷⁴

Essential Fatty Acids

Many BPH patients have received benefit from the administration of essential fatty acids (EFA) containing linoleic, linolenic, and arachidonic acids.³⁶ For reasons stated earlier, this could have implications for prostate cancer prevention or following conventional treatment for prostate cancer such as prostatectomy. An uncontrolled study of 19 subjects found reduction in residual urine in 12 of the subjects after several weeks of treatment.³⁷ This study confirmed other research which suggests EFA deficiencies in BHP and prostate cancer patients.³⁸ The combination of linoleic, linolenic, and arachidonic acids appears to be a critical factor in producing beneficial therapeutic effects.¹⁰⁸ A study found that severity of disease progression in a group of 49 prostate cancer patients was associated with a lower ratio of polyunsaturated-

to saturated fats in prostate tissue.¹⁷⁵ The polyunsaturated fats found in the study included linolenic acid, omega-3, and omega-6. A different study found that two commercial preparations of conjugated linoleic acid had inhibitory effects on the proliferation of prostate cancer cells.¹⁷⁶

Corn Oil

Evidence is emerging that the ingestion of large quantities of corn oil may increase the risk of developing prostate cancer and stimulating the growth of prostate cancer cells once developed. A study in mice found that corn oil and linoleic acid stimulated the growth of prostate cancer cells called DU145.¹²⁷ Rats fed a diet rich in corn oil (20% of fat intake) were more likely to have fast-growing prostate cancer cells than those on a fat-free diet.¹⁰⁹ An additional study found that rats fed a diet rich in fats obtained from corn oil (20%) were significantly more likely to develop carcinoma of the prostate than those fed a low-fat diet with low levels (5%) of corn oil.¹¹⁰ An additional study in mice using human prostate cancer cells that had been placed in the animals found that a diet rich (18%) in corn oil stimulated the growth of the cancerous cells to a greater extent than in mice who were fed a diet low in corn oil (5%).¹¹¹ A more recent study found that corn oil did not promote the development of prostate cancer in rats.¹⁷⁷

Amino Acids

A combination of amino acids that includes glutamic acid, alanine, and glycine given in the form of two 6 grain capsules three times daily for 2 weeks followed by one capsule three times daily has led to BPH symptom improvement in several studies.³⁹ A controlled study of 45 men with BPH found that delayed micturition occurred in 70% of cases, urine frequency was reduced in 73% of cases, nocturia was improved in 95% of cases, and urgency was reduced in 81% of cases.⁴⁰

Cholesterol

Cholesterol is an important factor in the formation of sexual hormones. Evidence indicates that cholesterol-lowering drugs have a beneficial effect on androgen formation in the prostate. A diet low in saturated fat coupled with regular aerobic exercise can reduce cholesterol levels in most individuals.

Soy

As previously noted, Asian populations have a reduced prevalence of progressive prostate cancer compared to Western populations. This may be due to generally high dietary soybean ingestion in these

populations. Soybeans have abundant quantities of phytoosterols. Phytoosterols are known to have cholesterol-lowering properties.⁴¹ Phytoosterols have demonstrated positive effects in treating BPH. A double-blind study evaluated the administration of 20 mg beta-sitosterol, a phytoesterol in soy, three times daily or placebo to 200 men with BPH.⁴² Beta-sitosterol increased maximum urinary flow rates from 9.9 ml/s to 15.2 ml/s and decreased mean residual urinary volume from 65.8 ml to 30.4 ml. The placebo group had no significant changes. Typical Western diets provide only about 80 mg/day of phytosterols compared to a traditional Japanese diet, which provides about 400 mg/day.⁴⁸ A three-and-a-half ounce serving of soybeans or tofu contains about 90 mg of beta-sitosterol. Higher consumption levels of soy and soyfoods has also been associated with a reduced risk of prostate cancer.⁴³ Two separate studies have attributed this effect to the isoflavonoids called daidzein and genistein.⁴⁴ Both of these agents have demonstrated 5-alpha-reductase reduction activity.

Saponins are another class of compound found in soybeans. Saponins are also found in many other types of plants. They are known to have a variety of anti-cancer properties.⁴⁵ Phytates are another compound found in soybeans and in other plants. Phytates have been shown to increase natural killer cell activity in the body.⁴⁶ The phytoestrogens found in soy are known to have anti-estrogenic effects.⁴⁷ Again, this can have important effects in the prevention of prostate cancer. It has been suggested that, in general, phytoestrogens have enzyme inhibition action, which could lead to inhibition of 5-alpha-reductase activity.⁴⁸

Asian cultures are believed to consume about 20-80 mg/day of genistein, whereas Western populations consume about 2-3 mg/day.⁴⁹ Messina and colleagues performed a review of 26 animal studies involving genistein and found that 17 (65%) of them demonstrated evidence of anti-carcinogenic activity from this compound.⁵⁰ It is believed that genistein is a natural PTK inhibitor.⁵¹ PTK inhibition has been found to be major factor in the inhibition of cancer.⁵² In vitro research has suggested genistein is a strong inhibitor of angiogenesis, a major factor in the growth of cancer cells.⁵³ Studies involving human volunteers who consumed soy beverages that contained 42 mg of genistein and 27 mg of daidzein daily produced peripheral blood concentrations of 0.5-1.0 mM.⁵⁴ This is known to be an insufficient quantity to inhibit the growth of cultured cancer

cells.⁵⁵ This suggests that these compounds may have a more important role as chemopreventative rather than chemotherapeutic agents.

Additional epidemiologic data has found that Japanese men have plasma levels of these isoflavone compounds that are 7 to 110 times higher than those in Finnish men.⁵⁶ The major constituent of these isoflavone concentrations is genistein. Researchers have suggested genistein may have a variety of mechanisms that prevent cancer formation, including direct growth inhibition, apoptosis induction, and cancer cell adhesion.⁵⁷ An in vitro study suggests that equol, biochanin A, and genistein are all potent inhibitors of 5-alpha-reductase.⁵⁸ Animal studies have suggested a diet high in genistein and daidzein resulted in a lower incidence of prostate cancer and longer disease-free periods after exposure to carcinogenic materials compared to a low isoflavone diet.⁵⁹ The isoflavones genistein and biochanin A inhibited many different human prostate cancer cell lines, but this effect did not occur with daidzein.⁶⁰

A new study found that 50 healthy men who received 50 mg of soy isoflavone twice daily for three weeks had an increase in antioxidant activity in prostate cells that could inhibit the formation of prostate cancer cells.¹¹² A different study found that the administration of genistein to prostate cancer cells already receiving radiation therapy increased the therapeutic effect of the therapy.¹¹³ An additional study has found that low doses of genistein can induce apoptosis in human prostate cancer cells.¹¹⁴

A more recent study in rats found that genistein in the diet decreased the incidence of poorly differentiated prostate tumors.¹⁷⁸ Researchers have also found that genistein regulates the expression of prostate-specific antigen, a key indicator of disease, in prostate cancer cells.¹⁷⁹ A different study found that genistein inhibited the proliferation of two prostate cancer cells lines (LNCaP and PC-3) by regulating an enzyme called glutathione peroxidase.¹⁸⁰ A recent study in mice found that genistein reduced the risk of developing prostate carcinoma in a dose-dependent manner.¹⁸¹ An additional study found that 50 mg of genistein twice daily for three weeks led to changes in healthy male subjects that are known to reduce the risk of developing prostate cancer.¹⁸² A randomized study of 35 Japanese men found that 400 ml daily consumption of soymilk led to significant decreases in serum levels of estrone, a hormone that may be an important factor in the development of prostate cancer.¹⁸³

Drug and Pesticide Avoidance

A variety of pesticides have been shown to produce changes that increase the activity of 5-alpha-reductase. Some of these compounds include dioxin, polyhalogenated biphenyls, hexachlorobenzene, diethylstilbestrol (DES), and dibenzofurans.⁶¹ A diet that avoids exposure to these agents is recommended for the prevention and treatment of prostate cancer.

Alcohol Avoidance

Beer ingestion has been associated with increased secretion of prolactin, which increases both testosterone formation and dihydrotestosterone conversion. Alcohol intake, in general, has negative effects on the prostate as evidenced by a 17-year study of 6,581 men in Hawaii. This study found that alcohol intake of at least 25 ounces per month was directly correlated with the development of BPH.⁶²

Saw Palmetto (*Serenoa repens*)

As mentioned earlier, there is no known connection between BPH and prostate cancer other than the fact that they have some similarities in etiology and symptoms. In both cases, the conversion of serum testosterone to DHT in the prostate gland appears to be an important part of the disease process. The drug finasteride has been used to treat BPH and is being studied in prostate cancer prevention trials. Finasteride works by inhibiting the conversion of testosterone to DHT. It accomplishes this by blocking the effects of 5-alpha-reductase, which converts testosterone to DHT.

Saw palmetto is the fat-soluble medicinal fraction obtained from the small palm tree called *Serenoa repens*, which grows in the West Indies and southeast Atlantic coast of North America.⁶³ Saw palmetto extract has been found to inhibit the conversion of testosterone to DHT in the prostate.⁶⁴ It has also demonstrated anti-estrogenic and receptor site-binding effects.⁶⁵ Surprisingly, estrogen has been found to contribute to the effects of BPH and possibly prostate cancer because it reduces the ability of the prostate to clear DHT levels.

A double-blind study of 35 men with BPH found that men given 160 mg of saw palmetto extract daily for 90 days had lower receptor values for estrogen and progesterone than those in a placebo group.⁶⁶ The researchers also found that saw palmetto extract blocks the translocation of the cytosol androgen receptor to the nucleus. The overall implications of the study are that saw palmetto extract has both anti-estrogenic and anti-androgenic effects. A double-blind, placebo-controlled study involving 110 BPH patients over 28

days found that saw palmetto administration significantly improved BPH symptoms, such as dysuria, nocturia, prostate volume, and residual urine.⁶⁷

Finasteride has had similar positive effects in the control of BPH but can take up to a year for the effects to occur, whereas saw palmetto generally produces positive effects in as little as 30 days and without serious adverse effects. Saw palmetto does not appear to effect PSA levels.

New research has discovered that men treated with saw palmetto had greater improvement in urinary tract symptoms associated with BPH than those who received placebo.¹¹⁵ A new randomized study of 44 men aged 45-80 years found that men receiving saw palmetto had improvement in symptoms associated with BPH, though this improvement did not reach statistical significance.¹¹⁶ A new microbiological study found that the application of an extract of saw palmetto to prostate cells led to increased rates of cell apoptosis.¹¹⁷

Researchers have found that saw palmetto has effects at the most basic physiologic levels by interfering with the effects of a hormone called prolactin and its stimulation of prostate cell proliferation.¹⁸⁴ A different group of researchers isolated a component of saw palmetto called myristoleic acid that induces cell death in the LNCaP line of prostate cancer cells.¹⁸⁵ Another study found that an extract of saw palmetto inhibits the activity of an enzyme that is important in the development of prostate cancer.¹⁸⁶ An extract of saw palmetto has been found to inhibit the activity of another enzyme called cyclooxygenase-2, which has been implicated in the progression of prostate cancer.¹⁸⁷ A recent study found that an extract of saw palmetto had equivalent therapeutic effects and fewer side effects as a commonly prescribed medication for prostate enlargement, a risk factor for development of prostate cancer.¹⁸⁸ Researchers found that saw palmetto has specific effects on mast cell accumulation and epithelium atrophy that may be responsible for its positive effects on prostatic hypertrophy and prostate cancer development.¹⁸⁹

Bitter Almond (*Pygeum Africanum*)

Pygeum africanum is an evergreen tree native to Africa. The fatty acids of the extract have similar properties to those of saw palmetto.⁶⁸ The tree also contains esters of ferulic acid, which have effects on the endocrine system. Studies have shown that docosanol reduces levels of testosterone and leutinizing

hormone.⁶⁹ Docosanol is a compound that has been found in *Pygeum africanum*. Docosanol has been found to reduce prolactin levels in the body.⁷⁰ Prolactin increases the uptake of testosterone and increases the conversion of testosterone to DHT in the body. Pygeum contains quantities of docosanol but the esters of ferulic acid have similar activity and are found in greater concentrations and are more bioavailable in pygeum than docosanol.⁷¹ Ferulic acid esters have been shown to reduce the cholesterol content in the prostate.⁷² Cholesterol end-products in the prostate have been associated with both BPH and cancer.⁷³ The sterolic portion of pygeum also works against the accumulation of testosterone in the prostate.⁷⁴

Double-blind studies have found that standardized pygeum extract, like that of saw palmetto, improved a variety of urinary symptoms associated with BPH, such as urinary frequency, nocturia, flow interruption, after-dribbling, weak stream, and hesitation.⁷⁵ Pygeum has also been found to increase prostatic secretions and improve the composition of the seminal fluid.⁷⁶ Standardized pygeum extract has also been found in double-blind clinical trials to improve the capacity of BPH patients to achieve erections.⁷⁷ All of these factors suggest a role for pygeum in prostate cancer patients who have received a prostatectomy, and possibly a role in the prevention of disease. A double-blind study found that saw palmetto was more effective than pygeum extract in most measures of BPH, but pygeum has better proven effects on prostate secretion.⁷⁸ The standardized extract of pygeum is 14% triterpenes including betasitosterol and 0.5% n-docosanol. The typical dose is 100-200 mg per day in divided doses. The crude herb is not used therapeutically.

New research in rat prostate tissues suggest pygeum africanum extract has positive effects on processes associated with BPH development.¹¹⁸ An additional study in animal tissues found further evidence of positive effects from pygeum africanum administration on inflammatory effects associated with the development of BPH.¹¹⁹ A review of 18 controlled clinical trials involving 1,562 men found that men treated with pygeum africanum for at least 30 days were more than twice as likely as controls to have symptom improvement associated with BPH than those receiving placebo.¹²⁰

Researchers have recently isolated an extract of pygeum africanum called V-1326 that may be responsible for its positive effects on prostate enlargement and the inhibition of prostate cancer

development.¹⁹⁰ A review of 18 randomized controlled trials found that *Pygeum africanum* is an effective treatment for an enlarged prostate and may have preventative effects on prostate cancer development.¹⁹¹ A study in rats found that an extract of *Pygeum africanum* called Tadenan counteracted many of the harmful physiologic effects of dihydrotestosterone.¹⁹² A different study involving rats found that Tadenan has a variety of physiologic effects that inhibit the proliferation of prostate cells.¹⁹³

Cernilton

Clinicians in Europe have used an extract of flower pollen called Cernilton for more than 35 years to treat prostatitis and BPH.⁷⁹ Double-blind clinical trials have shown that Cernilton improved BPH patients at an overall success rate of 70%.⁸⁰ Nocturia and diurnal frequency symptoms improve in about 70% of patients.⁸¹ Researchers have found evidence that Cernilton contains a substance that can inhibit the growth of prostate cells.⁸³ A study of 79 males aged 62-89 evaluated the effects of the administration of 63 mg Cernilton pollen extract daily for 12 weeks.⁸⁴ The mean baseline prostatic volume was 33.2 cm. During the study period, maximum urinary flow rate increased from 5.1 to 6.0 ml/sec. Average flow rates increased from 9.3 to 11 ml/sec. In addition, residual urinary flow rates decreased from 54.2 ml to below 30 ml. Furthermore, all of the following measures improved in these patients: urgency, intermittency, delayed voiding, post-void dribbling, prolonged voiding, incomplete emptying, dysuria, and nocturia. Cernilton demonstrated 50% inhibition on DU145 cell line growth after 2 days exposure at 5 mcg/ml.⁸⁵ A recent review of studies involving cernilton for the treatment of BPH found that cernilton modestly improves urologic symptoms associated with BPH.¹²¹

A review of studies involving cernilton and its effects on prostate enlargement found that cernilton improved urinary symptoms compared to a group receiving Tadenan and a control group in men with this condition.¹⁹⁴ A study of an extract of cernilton called DIBOA found that this compound inhibited the growth of DU-145 prostate cancer cells.¹⁹⁵ An additional study confirmed the inhibitory effects of DIBOA on a prostate cancer cell line.¹⁹⁶ A randomized study of 243 patients with urinary disturbances associated with benign prostatic hyperplasia found that a combination of tamsulosin hydrochloride with cernilton led to significant improvement in urinary symptoms.¹⁹⁷ Urinary difficulties are also a significant problem in prostate cancer patients. A study has also found that

cernilton pollen extract protects the epithelial cells lining the prostate, inhibits the proliferation of prostate cells, and enhances apoptosis of abnormal cells.¹⁹⁸

***Urtica dioica* (stinging needles)**

Urtica dioica has also been found to have beneficial effects in the treatment of BPH. Two double-blind studies have shown it to have greater efficacy in treating BPH than placebo.⁸⁶ The range of efficacy in this plant appears to be comparable to that of pygeum. This ranks it lower than saw palmetto for the treatment of BPH and in prostate cancer prevention potential.⁸⁷ The effects of *Urtica dioica* are similar to those of saw palmetto in that its main activity appears to be the interference of dihydrotestosterone binding to both cystolic and nuclear receptors.⁸⁸ This suggests a role in the prevention of prostate cancer. A study of 431 patients with BPH found that a combination therapy composed of extracts from saw palmetto and *urtica dioica* was as effective as finasteride in treating the symptoms of disease.¹²² A recent study found that a 20% extract of *Urtica dioica* had significant antiproliferative effects on human prostatic epithelial and stromal cells.¹⁹⁹

Garlic

A compound in aged garlic significantly inhibited the growth of human prostate cancer cells in vitro.⁹² A new study has demonstrated that garlic has properties that can detoxify cancer-producing factors.¹²³ Preliminary studies with allium, a component of garlic, suggest this compound has strong anti-tumor properties in both prostate and breast cancer cells.¹²⁴ A different study found evidence that a constituent of garlic called S-allylmercaptocyste has anti-tumor effects in prostate cells by converting the breakdown products of testosterone into less harmful substances.¹²⁵ A review article has cited new evidence for the antiproliferative effects of garlic on prostate cancer.²⁰⁰ An additional study found further evidence of the ability of garlic-derived allium to inhibit the proliferation of the LNCaP prostate cancer cell line.²⁰¹ A case-control study of 328 men diagnosed with prostate cancer found that men who consumed two or more servings of garlic per week had a 44% decreased risk of developing prostate cancer compared to those who never ate garlic.²⁰²

Citrus Pectin

Citrus pectin is a soluble component of plant fiber that is derived from citrus fruit. There is some evidence that citrus pectin can inhibit prostate cell formation and growth. A study in rats found that animals given 1.0% by volume citrus pectin in their

drinking water had statistically significant reductions in the metastasis of prostate carcinoma compared to control animals.¹³² A study involving mice found that the administration of high-dose (1.6 mg/ml) modified citrus pectin inhibited the tumor size after 20 days compared to controls.¹³³ A study of cancerous human prostate cells found that the addition of modified citrus pectin to the culture interfered with the mechanisms necessary for the cancer cells to multiply.¹³⁴ There is a significant amount of interest in studying modified citrus pectin for its anti-metastatic effects.¹³⁵ A newer study found evidence for the first time that modified citrus pectin can reduce the growth of solid primary tumors.²⁰³

Green Tea

Green tea has been shown to have inhibitory effects on prostate cancer cell growth. Studies in rats suggest that compounds contained in green tea inhibit the activity of 5-alpha-reductase, the enzyme that converts testosterone to dihydrotestosterone, which has carcinogenic effects in the prostate.¹³⁶ Researchers have found that the most potent of these compounds is called epigallocatechin-3-gallate (EGCG).¹³⁷ A study in rats found that green tea has inhibitory effects on other enzymes associated with the growth of prostate cancer cells.¹³⁸ EGCG has been found in high concentrations in the serum of green tea drinkers. EGCG and other compounds in green tea inhibit the activity of the enzyme called proteasome, a key factor in the formation of prostate cancer.¹³⁹

A recent review cited evidence that green tea can inhibit the formation of prostate cancer.²⁰⁴ A study in rats found that polyphenone-60, an extract of green tea, interfered with an enzyme called aromatase, which is known to be a key factor in the development of prostate cancer.²⁰⁵ Epidemiological evidence suggests green tea reduces the risk of developing prostate cancer, and a new study suggests green tea provides these protective effects by depleting levels of polyamines, which are an important part of cellular proliferation.²⁰⁶ A different study found evidence that green tea catechins inhibit some of the conversion of testosterone to dihydrotestosterone.²⁰⁷ A recent study in mice found that a polyphenolic fraction isolated from green tea not only inhibited localized prostate cancer growth but also inhibited the metastasis of the cancer to distant sites.²⁰⁸ A different study found that several catechins from green tea suppressed the growth of prostate cancer DU-145 cells.²⁰⁹

Spices

Spices add flavor to food but also may have cancer-fighting properties. India has one of the lowest cancer rates in the world, and the people there typically eat a diet containing a wide variety of spices. There is some limited evidence that capsaicin, the component of chile peppers that makes them hot, has some anti-cancerous properties. Both curry and cumin contain turmeric, which has been found to have anti-carcinogenic properties in cell cultures.¹⁴⁰ Research from animal studies in Asia has suggested capsaicin has anti-tumor effects.¹⁴³ A study at Yale University found that capsaicin interferes with the tumor-formation process in mice.¹⁴⁴

Curcumin

In studies involving mice, curcumin has demonstrated anti-carcinogenic activity in a variety of cancers.¹⁴¹ While the mechanisms of its action are still unknown, curcumin has also demonstrated anti-carcinogenic effects in human cancer cell lines.¹⁴² A recent study in mice found that curcumin decreased the proliferation of both androgen-dependent and androgen-independent prostate cancer cells.²¹⁰ A different study found that curcumin inhibits the growth of prostate cancer cells by inhibiting the enzymatic activity of tyrosine kinase at the epidermal growth factor receptor.²¹¹ A study determined that curcumin has great potential in the treatment of androgen-independent prostate cancer.²¹² A recent study found that curcumin inhibited both androgen-dependent and androgen-independent prostate cancer cell lines, and accomplishes this by inhibiting the effects of factors called NF-kappaB and AP-1.²¹³

Other Agents

A commercial eight-herb formulation called PC-SPES fed to rats in 0.05% and 0.025% levels in the diet led to dose-dependent effects on both tumor incidence and tumor growth rate.⁸⁹ PC-SPES decreased serum testosterone in six men with prostate cancer. An additional eight of eight patients had reduced PSA levels after using PC-SPES. PC-SPES has strong estrogenic activity that could cause adverse effects.⁹⁰ A study among 16 men with hormone-refractory prostate cancer led to significant improvement in quality-of-life measures, reduction in patient pain ratings, and decreased PSA levels.⁹¹

Biofeedback therapies can help men with prostate cancer and prostatectomy restore pelvic muscle function.⁹³ A treatment group regained continence after 51 days compared to 56 days in the control group.

Acupuncture was used on 7 men who developed hot flushes after receiving castration therapy for prostate cancer.⁹⁴ Six of the men completed 10 weeks of acupuncture therapy and had an average symptom decrease of 70%. A new study of ginseng found that a constituent called ginsenoside Rg3 demonstrated inhibitory action against a human prostate carcinoma cell line.¹²⁶ Evidence suggests licorice root (*Glycyrrhiza glabra*) can decrease circulating levels of testosterone in men. A study has found that licorice root has many similar properties to that of paclitaxel, a commonly used chemotherapeutic drug. This study also provided evidence that licorice root can induce apoptosis in prostate cancer cells.²¹⁴

Leading Hospitals for Urology

Below is the list of the 10 best hospitals in the U.S. for treatment of prostate cancer. This list was compiled from a survey of over 2500 doctors who were asked to pick the top hospitals for prostate cancer. *U.S. News and World Report* publishes this list every year.

1. Johns Hopkins Hospital
Baltimore, MD 21287
410-955-5464
2. Cleveland Clinic
Cleveland, OH 44195
800-223-2273, 216-444-2200
3. Mayo Clinic
Rochester, MN 55905
507-284-2511
4. UCLA Medical Center
Los Angeles, CA 90095
310-825-9111
5. New York Presbyterian Hospital
New York, NY 10032
212-305-2500
6. Duke University Medical Center
Durham, NC 27710
919-684-8111
7. Stanford University Hospital
Stanford, CA 94305
415-723-4000
8. Massachusetts General Hospital
Boston, MA 02114
617-726-2000

9. Memorial Sloan-Kettering Cancer Center
New York, NY 10021
212-639-2000

10. Barnes-Jewish Hospital
St. Louis, MO 63110
314-362-5000

Complimentary and Alternative Physicians Directories

To locate a physician in your area that practices complimentary and alternative medicine, visit these web sites.

Alternative Medicine Directory

www.altmedicine.net

Includes directory for Alternative Medicine Doctors who specialize in different fields according to a particular ailment.

The New World of Holistic Health

www.earthmed.com

Free membership. Selection of over 11,000 earthmed.com practitioners depending on state and type of treatment.

Holistic Health Yellow Pages

www.findhealer.com

Enrollment necessary. Includes directory of doctors according to ailment.

American Association of Naturopathic Physicians

www.naturopathic.org

Headed up by Andrew L. Rubman and Susan Gordon. Search ability to find an ND according to area or name if you know the name of the doctor.

Health USA

www.healthusa.com

Under related searches this web site gives many options including: Alternative Medicine here you can find a lot of information, and Doctors which shows where Doctors are located, ability to ask questions via email and more.

Lori's Natural Food Center

www.lorisnatural.com

Under the category About US and the subcategory Doctor there is a listing of doctors from New York and Massachusetts.

American College for the Advancement in Medicine

www.acam.org

ACAM is committed to the education of physicians on nutritional and preventive medicine and alternative procedures. A search is available to find doctors both by locality or by name.

Ask Dr. Weil

www.drweil.com

The web site listing for Dr. Weil provides the opportunity to ask him questions. Under the search engine on this site find other Alternative Doctors locally.

American Academy of Medical Acupuncture (AAMA)

<http://medicalacupuncture.org>

820 Wilshire Boulevard, Suite 500

Los Angeles, California 90036

Tel: (323) 937-5514

Email: JDOWDEN@prodigy.net

Organization of physician acupuncturists. Medical acupuncture is the term used to describe acupuncture performed by a doctor trained and licensed in Western medicine who has also had thorough training in acupuncture as a specialty practice. Such a doctor can use one or the other approach, or a combination of both as the need arises, to treat an illness.

American Association of Naturopathic Physicians

<http://www.naturopathic.org/>

601 Valley Street, Ste. 105.

Seattle, WA 98109

Tel: (206) 298-0126

Fax: (206) 298-0129

Email: webmaster@naturopathic.org

This organization represents the naturopathic physicians from the four recognized schools of naturopathy - see Herbal Medicine Resource Guide. The web site contains a physician referral listing, searchable databases of information and resources on natural healing by licensed Naturopathic Doctors.

Society for Nutritional Oncology

www.noat.org

3455 Salt Creek Lane

Arlington Heights, IL 60005

email: NOATSociety@noat.org

Footnotes

- 1 Hayes RB et al. Dietary factors and risks for prostate cancer among blacks and whites in the United States. *Cancer Epidemiol Biomarkers Prev* 1999; 8 (1):25-34.
- 2 Chyou PH et al. A prospective study of alcohol, diet, and other lifestyle factors in relation to obstructive uropathy. *Prostate* 1993;22:253-264.
- 3 DeRosa G et al. Prolactin secretion after beer. *Lancet* 1981;2:934.
- 4 Adlercreutz H et al. Phyto-oestrogens and Western diseases. *Ann Med* 1997;29:95-120.
- 5 Grant WB et al. An ecologic study of dietary links to prostate cancer. *Altern Med Rev* 1999 Jun;4:162-169
- 6 Bendich A et al. Biological action of carotenoids. *FASEB J* 1989;3:1927-1932.
- 7 Francheschi S et al. Tomatoes and risk of digestive-tract cancers. *Int J Cancer* 1994;59:181-184.
- 8 Gann PH et al. Lower prostate cancer risk in men with elevated plasma lycopene levels; results of a prospective analysis. *Cancer Res* 1999 Mar 15;59:1225-1230.
- 9 Rao AV et al. Serum and tissue lycopene and biomarkers of oxidation in prostate cancer patients: a case-control study. *Nutr Cancer* 1999;33:159-164.
- 10 Norrish AE et al. Prostate cancer and dietary carotenoids. *Am J Epidemiol* 2000;151:119-123.
- 11 Giovannucci E et al. Intake of carotenoids and retinal in relation to risk of prostate cancer. 1995; 87:1767-1776.
- 12 Combs et al. Reduction of cancer risk with an oral supplement of selenium. 1997; 10:227-234.
- 13 Webber et al. Inhibitory effects of selenium on the growth of DU-145 human prostate carcinoma cell in vitro. *Biochem Biophys Res Commun* 1985; 130:603-609.
- 14 Hardell L et al. Levels of selenium in plasma and glutathione peroxidase in erythrocytes in patients with prostate cancer or benign hyperplasia. *Eur J Cancer Prev* 1995;4:91-95.
- 15 Webber MM. Selenium prevents the growth stimulatory effects of cadmium on human prostatic epithelium. *Biochem Biophys Res Commun* 1985;127:871-877.
- 16 Yoshizawa K et al. Study of prediagnostic selenium level in toenails and the risk of advanced prostate cancer. *J Natl Cancer Inst* 1998;90:1219-1224.
- 17 Nomura AM et al. Serum selenium and subsequent risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2000;9:883-887.
- 18 Willett WC et al. Prediagnostic serum selenium and risk of cancer. 1983;2:130-134.
- 19 Combs GF et al. Reduction of cancer risk with an oral supplement of selenium. *Biomed Environ Sci* 1997;10:227-234.
- 20 Clark LC et al. Decreased incidence of prostate cancer with selenium supplementation: results of a double-blind cancer prevention trial. *Br J urol* 1998;81:730-734.
- 21 Hartman TJ et al. The association between baseline vitamin E, selenium, and prostate cancer in the alpha-tocopherol, beta-carotene cancer prevention study. *Cancer Epidemiol Biomarkers Prev* 1998;7: 335-340.
- 22 Pastori M et al. Lycopene in association with alpha-tocopherol inhibits at physiological concentrations proliferation of prostate carcinoma cells. *Biochem Biophys Res Commun* 1998;250:582-585.
- 23 Pollar M. Vitamin D analog suppressed tumor development in rats. *Prostate* 1999;39:305-309.
- 24 Loseshwar BL et al. Inhibition of prostate cancer metastasis in vivo: a comparison of 1,25-dihydrovitamin D (calcitrol) and EB 1089
- 25 Blutt SE et al. Vitamin D and prostate cancer. *Proc Soc Exp Biol Med* 1999;221:89-98.
- 26 Schwarz GG et al. 1alpha,25-Dihydroxyvitamin D (calcitrol) inhibits the invasiveness of human prostate cancer cells. 1997 *Cancer Epidemiol Biomarkers Prev* 1997; 6:727-732.
- 27 Miller GJ. Vitamin D and prostate cancer: biologic interactions and clinical potentials 1998-99;17:353-360.
- 28 Bush IM et al. Zinc and the prostate. Presented at the annual meeting of the AMA,2974.
- 29 Fahim M et al. Zinc treatment for the reduction of hyperplasia of the prostate. *Fed Proc* 1976;35:361.
- 30 Leake A et al. Subcellular distribution of zinc in the benign and malignant human prostate: evidence for a direct zinc androgen interaction. *Acta Endocrinol* 1984;105:281-288.
- 31 Zaichick et al. Zinc concentration in human prostatic fluid. Normal, chronic prostatitis, adenoma, and cancer. *Int Urol Nephrol* 1996;28:687-694.
- 32 Leake A et al. Subcellular distribution of zinc in the benign and malignant human prostate: evidence for a direct zinc androgen interaction. *Acta Endocrinol* 1984;105:281-288.
- 33 Login IS et al. Zinc may have a physiological role in regulating pituitary prolactin secretion. *Neuroendocrinology* 1983;37:317-320.
- 34 Farnsworth WE et al. Interaction of prolactin and testosterone in the human prostate. *Urol Res* 1981;9:79-88.

- 35 Judd AM et al. Zinc acutely, selectively, and reversibly inhibits pituitary prolactin secretion. *Brain Res* 1984;294:190-192.
- 36 Hart JP et al. Vitamin F in the treatment of prostatic hyperplasia. Report Number 1. Milwaukee, WI: Lee Foundation for Nutritional Research. 1941
- 37 Scott, WW. The lipids of the prostatic fluid, seminal plasma, and enlarged prostate gland of man. *J Urol* 1945; 53:712-718.
- 38 Boyd EM et al. Prostatic hypertrophy as part of a generalized metabolic disease. Evidence of the presence of a lipopenia. *J Urol* 1939;41:406-411.
- 39 Dumrau F. Benign prostatic hyperplasia: amino acid therapy for symptomatic relief. *Am J Ger* 1962; 10:426-430.
- 40 Feinblatt HM et al. Palliative treatment of benign prostatic hypertrophy: value of glycine, alanine, glutamic acid combination. *J Maine Med Assoc* 1958; 49:99-102.
- 41 Tilvis RS et al. Serum plant sterols and their relation to cholesterol absorption. *Am J Clin Nutr* 1986; 43:92-97.
- 42 Berges RR et al. Randomized, place-controlled, double-blind clinical trial of beta-sitosterol in patients with benign prostatic hyperplasia. *Lancet* 1995;345:1529-1532.
- 43 Morton MS et al. The preventive role of diet in prostatic disease. *Br J Urol* 1996; 77:481-493.
- 44 Messina M et al. The roles of soy products in reducing the risk of cancer. *J Natl Cancer Inst*
- 45 Elias R et al. Antimutagenic activity of some saponins isolated from *Calendula officinalis* L., *C. arvensis* L., and *Hedera helix* L. *Mutagenesis* 1990; 5:327-331.
- 46 Baten A et al. Inositol-phosphate-induced enhanced of natural killer cell activity correlates with tumor suppression. *Carcinogenic* 1989; 10:1595-1598.
- 47 Adlercreutz H et al. Phyto-oestrogens and Western diseases. *Ann Med* 1997;29:95-120.
- 48 Evans BA et al. Inhibition of 5 alpha-reductase in genital skin fibroblasts and prostate tissue by dietary lignans and isoflavonoids. *J Endocrinol* 1995; 147:295-302.
- 49 Messina M et al. The role of soy products in reducing the risk of cancer. *J Natl Cancer Inst* 1991;83:541-546.
- 50 Messina M et al. Soy intake and cancer risk: a review of the in vitro and in vivo data. *Nutr Cancer* 1994; 21:113-131.
- 51 Barnes S et al. Rationale for the use of genistein-containing soy matrices in chemoprevention trials for breast and prostate cancer. *J Cell Biochem Suppl* 1995; 22:181-187.
- 52 St Clair M et al. Suppression of dimethylhydrazine-induced carcinogenesis in mice by dietary addition of the Bowman-Birk Protease inhibitor. *Cancer Res* 1990;50:580-586.
- 53 Fotsis T et al. Genistein, a dietary-derived inhibitor of in vitro angiogenesis. *Proc Natl Acad Sci USA* 1993; 90: 2690-2694.
- 54 Barnes S et al. Soy isoflavonoids and cancer prevention. Underlying biochemical and pharmacological issues. *Adv Exp Med Biol* 1996; 401:87-100.
- 55 Barnes S et al. Biochemical targets of the isoflavone genistein in tumor cell lines. *Proc Soc Exp Biol Med* 1995; 208:103-108.
- 56 Adlercreutz H et al. Plasma concentrations of phyto-oestrogens in Japanese men. *Lancet* 1993; 342:1209-1210.
- 57 Kyle E et al. Genistein-induced apoptosis of prostate cancer cells is preceded by a specific decrease in focal adhesion kinase activity. *Mol Pharmacol* 1997; 51:193-200.
- 58 Evans BA et al. Inhibition of 5 alpha-reductase in genital skin fibroblasts and prostate tissue by dietary lignans and isoflavonoids. *J Endocrinol* 1995; 147:295-302.
- 59 Pollard M et al. Influence of isoflavones in soy protein isolates on development of induced prostate-related cancers in L-W rats. *Nutr Cancer* 1997; 28:41-45.
- 60 Peterson G et al. Genistein and biochanin A inhibit the growth of human prostate cancer cells but not epidermal growth factor receptor tyrosine autophosphorylation. *Prostate* 1993; 22:335-345.
- 61 Kappas A et al. Nutrition-endocrine interactions. Induction of reciprocal changes in the delta-5-alpha-reduction of testosterone and the cytochrome P-450-dependent oxidation of estradiol by dietary macronutrients in man. *Proc Natl Acad Sci USA* 1983; 80: 7646-7649.
- 62 Chyou PH et al. A prospective study of alcohol, diet, and other lifestyle factors in relation to obstructive uropathy. *Prostate* 1993; 22:253-264.
- 63 Duke JA. Handbook of medicinal herbs. Boca Raton, FL: CRC Press. 1985:p118.
- 64 Carilla E et al. Binding of Permixon, a new treatment for prostatic benign hyperplasia to the cytosolic androgen receptor in the rat prostate. *J Steroid Biochem* 1984; 20: 521-523.
- 65 Sultan C et al. Inhibition of androgen metabolism and binding by a liposteric extract of *Serenoa repens* B in human foreskin fibroblasts. *J Steroid Biochem* 1984; 20:515-519.

- 66 Carilla E et al. Binding of Permixon, a new treatment for prostatic benign hyperplasia to the cytosolic androgen receptor in the rat prostate. *J Steroid Biochem* 1984; 20: 521-523.
- 67 Crimi A et al. Extract of *Serenoa repens* for the treatment of the functional disturbances of prostate hypertrophy. *Med Praxis* 1983; 4: 47-51.
- 68 Hinman F. *Benign Prostatic Hyperplasia*. New York: Springer-Verlag. 1983
- 69 Martinelli EM et al. Characterization of *Pygeum africanum* bark extracts by HRCG with computer assistance, *HRC & CC* 1986; 9:106-110.
- 70 Hinman F. *Benign Prostatic Hyperplasia*. New York: Springer-Verlag. 1983.
- 71 Uberti E. HPLC analysis of n-docosyl ferulate in *Pygeum africanum* extracts and pharmaceutical formulations. *Fitotherapia* 1990; 41: 342-347.
- 72 Bombardelli E. Methods, composition and compounds for the treatment of prostatic adenoma. *EP Appl* 8330491.3, June 10, 1985.
- 73 Hinman F. *Benign Prostatic Hyperplasia*. New York: Springer-Verlag. 1983.
- 74 Marcoli M. Anti-inflammatory and antiedemigenic activity of extract of *Pygeum africanum* in the rat. *New Trends Androl Sci* 1985; 1:89.
- 75 Colpi G et al. Study of the activity of chloroformic extract of *Pygeum africanum* bark in the treatment of urethral obstructive syndrome caused by non-cancerous prostatic hyperplasia. *Urologia* 1976; 43:441-448.
- 76 Luchetta G et al. Reactivation from the prostatic gland in cases of reduced fertility. *Urol Int* 1984; 39: 222-224.
- 77 Carani C et al. Urological and sexual evaluation of treatment of benign prostatic disease using *Pygeum africanum* at high dose. *Arch Ital Urol Nefrol Androl* 1991; 63: 341-345.
- 78 Duvia R et al. Advances in the phytotherapy of prostatic hypertrophy. *Med Praxis* 1983; 4:143-148.
- 79 Yasumoto R et al. Clinical evaluation of long-term treatment using Cernilton pollen extract in patients with benign prostatic hyperplasia. *Clinical Therapeutics* 1995; 17:82-86.
- 80 Buck AC et al. Treatment of outflow tract obstruction due to benign prostatic hyperplasia with the pollen extract Cernilton. A double-blind, placebo-controlled study. *Br J Urol* 1990; 66:398-404.
- 81 Dutkiewicz S. Usefulness of Cernilton in the treatment of benign prostatic hyperplasia. *Int Urol Nephrol* 1996; 28: 49-53
- 82 Habib FK et al. Identification of a prostate inhibitory substance in a pollen extract. *The Prostate* 1995; 26:133-139.
- 83 Yasumoto R et al. Clinical evaluation of long-term treatment using Cernilton pollen extract in patients with benign prostatic hyperplasia. *Clinical Therapeutics* 1995; 17:82-86.
- 84 Habib FK et al. Identification of a prostate inhibitory substance in a pollen extract. *Prostate* 1995; 26:133-139.
- 85 Belaiche P et al. Clinical studies on the palliative treatment of prostatic adenoma with extract of *Urtica* root. *Phytother res* 1991; 5:267-269.
- 86 Romics I et al. Observations with Bazoton in the management of prostatic hyperplasia. *Int Urol Nephrol* 1987; 19: 293-297.
- 87 Wagner H. Search for the antiprostatic principle of stinging nettle (*Urtica dioica*) roots. *Phytomedicine* 1994; 1:213-224.
- 88 Raloff J et al. Radical prostates. *Science News* 1997 151: 126-27.
- 89 Tiwari RK et al. Anti-tumor effects of PC-SPES, an herbal formulation in prostate cancer. 1999; 14:713-719.
- 90 Geliebter J et al. PC-SPES in prostate cancer. *New England Journal of Medicine* 1999; 340: 567-568.
- 91 Pfeifer BL et al. PC-SPES, a dietary supplement for the treatment of hormone-refractory cancer. *BJU Int* 2000. 85:481-485.
- 92 Mathewson-Chapman M. Pelvic muscle exercise/biofeedback for urinary incontinence after prostatectomy. *J Cancer Educ* 1997; 12: 218-23.
- 93 Hammar M et al. Acupuncture treatment of vasomotor symptoms in men with prostatic carcinoma: a pilot study. *J Urol* 1999; 161: 853-56.
- 94 Giovannuci E, et al. Intake of carotenoids and retinal in relation to risk of prostate cancer. *J Nat Cancer Inst.* 1995; 87: 1767-76.
- 95 Guttenplan JB, et al. Effects of a lycopene-rich diet on spontaneous and benzo[a] pyrene-induced mutagenesis in prostate, colon, and lungs of the lacZ mouse. *Cancer Lett* 2001; 164:1-6.
- 96 Agarwal S, Rao AV. Tomato lycopene and its role in human health and chronic diseases. *CMAJ* 2000; 163:739-44.
- 97 Jiang C, et al. caspases as key executors of methyl selenium-induced apoptosis (anoikis) of DU-145 prostate cancer cells. *Cancer Res* 2001; 61:3062-3070.
- 98 Hartman TJ, et al. Effects of long-term alpha-tocopherol supplementation on serum hormones in older men. *Prostate* 2001; 46: 33-38.
- 99 Helzlsouer KJ, et al. Association between alpha-tocopherol, gamma-tocopherol, selenium, and subsequent prostate cancer. *J Natl Cancer Inst* 2000; 92: 1966-67.

- 101 Gunawardena K, et al. Vitamin E and other antioxidants inhibit human prostate cancer cells through apoptosis. *Prostate* 2000; 44: 287-95.
- 102 Hsu JY, Feldman D, McNeal JE, Peehl DM. Reduced 1 alpha-hydroxylase activity in human prostate cancer cells correlates with decreased susceptibility to 25-hydroxyvitamin D3-induced growth inhibition. *Cancer Res* 2001;61;2852-6.
- 103 Rashid SF, et al. Synergistic growth inhibition of prostate cancer cells by 1 alpha Dihydroxyvitamin D(3) and its 19-nor-hexafluoride analogs in combination with either sodium butyrate or trichostatin A. *Oncogene* 2001; 20: 1860-72.
- 104 Zhao XY, Feldman D. The role of vitamin D in prostate cancer. *Steroid* 2001; 66: 293-300.
- 105 Yu CL, Tsai MH. Fetal fetuin selectively induces apoptosis in cancer cell lines and shows anti-cancer activity in tumor animal models. *Cancer Lett* 2001;166: 173-84.
- 106 Ishii K, et al. Aminopeptidase N regulated by zinc in human prostate participates in tumor cell invasion. 2001; 92: 49-54.
- 107 Ishii K, et al. Inhibition of aminopeptidase N (AP-N) and urokinase-type plasminogen activator (uPA) by zinc suppresses the invasion activity in human urological cancer cells. *Biol Pharm Bull* 2001; 24: 226-30.
- 108 Ghosh J, Myers CE. Arachidonic acid stimulates prostate cancer cell growth: critical role of 5-lipoxygenase. *Biochem Biophys Res Commun* 1997; 235: 418-23.
- 109 Clinton SK, et al. Growth of Dunning transplantable prostate adenocarcinomas in rats fed diets with various fat contents. *J Nutr* 1988; 118: 908-914.
- 110 Kondo Y, et al. Promotional effect of two-generation exposure to a high-fat diet prostate carcinogenesis in ACI/Seg rats. *Cancer Res* 1994; 54: 6129-32.
- 111 Connolly JM, et al. Effects of dietary fatty acids on DU145 human prostate cancer cell growth in athymic nude mice. *Nutr Cancer* 1997; 29: 114-9.
- 112 Davis JN, et al. Soy isoflavone supplementation in healthy men prevents NF-kappaB activation by TNF-alpha in blood lymphocytes. *Free Radic Biol Med* 2001;30:1293-1302.
- 113 Hillman GG, et al. Genistein potentiates the radiation effect of prostate carcinoma. *Clin Cancer Res* 2001; 7: 382-90.
- 114 Shen JC, et al. Low-dose genistein cyclin-dependent kinase inhibitors a G(1) cell-cycle arrest in human prostate cancer cells. *Mol Carcinog* 2000; 29:92-102.\
- 115 Wilt TJ, et al. Phytotherapy for benign prostatic hyperplasia. *Public Health Nutr* 2000; 3: 459-72.
- 116 Marks LS, et al. Effects of a saw palmetto herbal blend in men with symptomatic benign prostatic hyperplasia. *J Urol* 2000; 163: 151-6.
- 117 Iguchi K, et al. Myristoleic acid, a cytotoxic component in the extract from *Serenoa repens*, induces apoptosis and necrosis in human prostate LNCaP cells. *Prostate* 2001; 47: 59-65.
- 118 Solano RM, et al. Effects of *Pygeum africanum* extract (Tadenan) on vasoactive intestinal peptide receptors, G proteins, and adenylyl cyclase in rat ventral prostate. *Prostate* 2000; 45: 245-52.
- 119 Szolnoki E, et al. The effect of *Pygeum africanum* on fibroblast growth factor (FGF) and transforming growth factor beta (TGF beta 1/LAP) expression in animal model. *Acta Microbiol Immunol Hung* 2001; 48: 1-9.
- 120 Ishani A, et al. *Pygeum africanum* for the treatment of patients with benign prostatic hyperplasia: a systematic review and quantitative meta-analysis. *Am J Med* 2000; 109: 654-64.
- 121 Wilt T, et al. Cernilton for benign prostatic hyperplasia. *Cochrane Database Syst Rev* 2000: CD001042.
- 122 Sokeland J. Combined sabal and urtica extract compared with finasteride in men with benign prostatic hyperplasia: analysis of prostate volume and therapeutic outcome. *BJU Int* 2000; 86: 439-42.
- 123 Lamm DL, Riggs DR. Enhanced immunocompetence by garlic: role in bladder cancer and other malignancies. *J Nutr* 2001; 131: 1067S-70S.
- 124 Pinto JT, Rivlin RS. Antiproliferative effects of allium derivatives from garlic. *J Nutr* 2001; 131: 1058S-60S.
- 125 Pinto, JT, et al. Alterations of prostate biomarker expression and testosterone utilization in huam LNCaP prosatic carcinoma cells by garlic-derived S-allylmercaptocysteine. *Prostate* 2000; 45: 304-14.
- 126 Liu WK, et al. Anti-proliferative effect of ginseng saponins on human prostate cancer cell line. *Life Sci* 2000; 67: 1297-1306.
- 127 Connolly JM, et al. Effects of dietary fatty acids on DU145 human prostate cancer cell growth in athymic nude mice. *Nutr Cancer* 1997; 29: 114-19.
- 128 Rodriguex C, et al. Body mass index, height, and prostate cancer mortality in two large cohorts of adult men in the United States. *Cancer Epidemiol Biomarkers* 2001; 10: 345-53.
- 129 Hsing AW, Deng J, Sesterhenn IA, Mostofi FK, Stanczyk FZ, Benichou J, Xie T, Gao YT. *Cancer Epidemiol Prev* 2000; 9: 1335-41.
- 130 Putnam SD, et al. Lifestyle and anthropometric risk factors for prostate cancer in a cohort of Iowa men. *Ann Epidemiol* 2000; 10: 361-9.

- 131 Giovannucci E, et al. Height, weight, and risk of prostate cancer. *Cancer Epidemiol Biomarkers* 1997; 6: 557-63.
- 132 Pienta KJ, et al. Inhibition of spontaneous metastasis in a rat prostate cancer model by oral administration of modified citrus pectin. *J Natl Cancer Inst* 1995; 87: 348-53.
- 133 Hayashi A, Gillen AC, Lott JR. Effects of daily oral administration of quercetin chalcone and modified citrus pectin. *Altern Med Rev* 2000; 5: 546-52.
- 134 Hsieh TC, Wu JM. Changes in cell growth, cyclin/kinase, endogenous phosphoproteins and nm23 gene expression in human prostatic JCA-1 cells treated with modified citrus pectin. *Biochem Mol Biol Int* 1995; 37: 833-41.
- 135 Modified citrus pectin-monograph. *Altern Med Rev* 2000; 5: 573-5.
- 136 Liao S, Hiipakka RA. Selective inhibition of steroid 5 alpha-reductase isozymes by tea epicatechin-3-gallate and epigallocatechin-3-gallate. *Biochem Biophys Res* 1995; 214: 833-38.
- 137 Paschka AG, Butler R, Young CY. Induction of apoptosis in prostate cancer cell lines by the green tea component, (-)-epigallocatechin-3-gallate. *Cancer Lett* 1998; 130: 1-7.
- 138 Gupta S, et al. Prostate cancer chemoprevention by green tea: in vitro and in vivo inhibition of testosterone-mediated induction of ornithine decarboxylase. *Cancer Res* 1999; 59: 2115-20.
- 139 Nam S, Smith DM, Dou QP. Ester bond-containing tea polyphenols potently inhibit proteasome activity in vitro and in vivo. *J Biol Chem* 2001; 276: 13322-30.
- 140 Rao CV, et al. Chemoprevention of colon carcinogenesis by dietary curcumin, a naturally occurring plant phenolic compound. *Cancer Res* 1995; 55: 259-66.
- 141 Huang MT, Newmark HL, Frenkel K. Inhibitory effects of curcumin on tumorigenesis in mice. *J Cell Biochem Suppl* 1997; 27: 26-34.
- 142 Hanif R, Qiao L, Shiff SJ, Rigas B. Curcumin, a natural plant phenolic food additive, inhibits cell proliferation and induces cell cycle changes in colon adenocarcinoma lines by a prostaglandin-independent pathway. *J Lab Clin Med* 1997; 130: 576-84.
- 143 Yun TK. Update from Asia. Asian studies on cancer chemoprevention. *Ann NY Acad Sci* 1999; 889: 157-92.
- 144 Surh YJ, et al. Chemoprotective effects of capsaicin and diallyl sulfide against mutagenesis or tumorigenesis by vinyl carbamate and N-nitrosodimethylamine. *Carcinogenesis* 1995; 16: 2467-71.
- 145 Zigler RG. Vegetables, fruits, and carotenoids and the risk of cancer. *Am J Clin Nutr* 1991; 53: 2515-2595.
- 146 Bhatia N, Agarwal R. Detrimental effect of cancer preventive phytochemicals silymarin, genistein, and epigallocatechin 3-gallate on epigenetic events in human prostate carcinoma DU145 cells. *Prostate* 2001; 46: 98-107.
- 147 Waladkhani AR, Clemens MR. Effect of dietary phytochemicals on cancer development. *Int J Mol Med* 1998; 1: 747-53.
- 148 Craig WJ. Phytochemicals: guardians of our health. *J Am Diet Assoc* 1997; 97: S199-204.
- 149 Reuter News Service. May 17, 2002.
- 150 De Marzo A, et al. *Cancer Research* 2002.
- 151 Van Breemen RB, et al. Liquid chromatography-mass spectrometry of cis- and all-trans-lycopene in human serum and prostate tissue after dietary supplementation with tomato sauce. *J Agric Food Chem* 2002;50:2214-9.
- 152 Fairfield KM, Fletcher RH. Vitamins for chronic disease prevention in adults: scientific review. *JAMA* 2002;287:3116-26.
- 153 Giovannuci E, et al. A prospective study of tomato products, lycopene, and prostate cancer risk. *J Natl Cancer Inst* 2002;94:391-8.
- 154 Bowen P, et al. *J Natl Cancer Inst* Dec 19, 2001.
- 155 BW HealthWire. July 13, 2001.
- 156 Moyad MA. Selenium and vitamin E supplements for prostate cancer: evidence or embellishment? *Urology* 2002;59:9-19.
- 157 Israel K, et al. Vitamin E succinate induces apoptosis in human prostate cancer cells: role for Fas in vitamin E succinate-triggered apoptosis. *Nutr Cancer* 2000;36:90-100.
- 158 Hartman TJ, et al. Effects of long-term alpha-tocopherol supplementation on serum hormones in older men. *Prostate* 2001;46:33-8.
- 159 Salih HR, et al. Retinoic acid and vitamin E modulate expression and release of CD178 in carcinoma cells: consequences for induction of apoptosis in CD95-sensitive cells. *Exp Cell Res* 2001;270:248-58.
- 160 Lagiou A, et al. Are there age-dependent effects of diet on prostate cancer risk? *Soz Praventivmed* 2001;46:329-34.
- 161 Moyad MA. Complementary/alternative therapies for reducing hot flashes in prostate cancer patients: reevaluating the existing indirect data from studies of breast cancer and postmenopausal women. *Urology* 2002;59:20-33.

- 162 Yeh S, et al. Proceedings of the National Academy of Sciences. May 28, 2002.
- 163 Finley JW, et al. Journal of Agricultural and Food Chemistry. 2001;49:2679-83.
- 164 Rashid SF, et al. Synergistic growth inhibition of prostate cancer cells by 1 alpha, 25 Dihydroxyvitamin D(3) and its 19-nor-hexafluoride analogs in combination with either sodium butyrate or trichostatin A. *Oncogene* 2001;20:1860-72.
- 165 Peehl DM, Seto E, Feldman D. Rationale for combination ketoconazole/vitamin D treatment of prostate cancer. *Urology* 2001;58:123-6.
- 166 Beer TM, et al. Weekly high-dose calcitriol and docetaxel in advanced prostate cancer. *Semin Oncol* 2001;28:49-55.
- 167 Crescioli C, et al. Vitamin D3 analogue inhibits keratinocyte growth factor signaling and induces apoptosis in human prostate cancer cells. *Prostate* 2002;50:15-26.
- 168 Polek TC, Weigel NL. Vitamin D and prostate cancer. *J Androl* 2002;23:9-17.
- 169 Yang ES, et al. Vitamin D-mediated growth inhibition of an androgen-ablated LNCaP cell line model of human prostate cancer. *Mol Cell Endocrinol* 2002;186:69-79.
- 170 Hsieh CL, et al. A novel targeting modality to enhance adenoviral replication by vitamin D(3) in androgen-independent human prostate cancer cells and tumors. *Cancer Res* 2002;62:3084-92.
- 171 Ishii K, et al. Inhibition of aminopeptidase N (AP-N) and urokinase-type plasminogen activator (uPA) by zinc suppresses the invasion activity in human urological cancer cells. *Biol Pharm Bull* 2001;24:226-30.
- 172 Yu CL, Tsai MH. Fetal fetuin induces apoptosis in cancer cell lines and shows anti-cancer activity in tumor animal models. *Cancer Lett* 2001;166:173-84.
- 173 Narla G, et al. KLF6, a candidate tumor suppressor gene mutated in prostate cancer. *Science* 2001;294:2563-6.
- 174 Bataineh ZM, et al. Zinc in normal and pathological human prostate gland. *Saudi Med J* 2002;23:218-20.
- 175 Freeman VL, et al. Prostatic levels of fatty acids and the histopathology of localized prostate cancer. 2000;164:2168-72.
- 176 Palombo JD, et al. The antiproliferative effects of biologically active isomers of conjugated linoleic acid on human colorectal and prostatic cancer cells. *Cancer Lett* 2002;177:163-72.
- 177 Mori T, et al. Beef tallow, but not perilla or corn oil, promotion of rat prostate and intestinal carcinogenesis by 3,2'-dimethyl-4-aminobiphenyl. *Jpn J Cancer Res* 2001;92:1026-33.
- 178 Lamartiniere CA, et al. Genistein chemoprevention: timing and mechanisms of action in murine mammary and prostate. *J Nutr* 2002;132:552S-558S.
- 179 Davis JN, Kucuk O, Sarkar FH. Expression of prostate-specific antigen is transcriptionally regulated by genistein in prostate cancer cells. *Mol Carcinog* 2002;34:91-101.
- 180 Suzuki K, et al. Genistein, a soy isoflavone, induces glutathione peroxidase in the human prostate cancer cell lines LNCaP and PC-3. *Int J Cancer* 2002;99:846-52.
- 181 Mentor-Marcel R, et al. Genistein in the diet reduces the incidence of poorly differentiated prostatic adenocarcinoma in transgenic mice (TRAMP). *Cancer Res* 2001;61:6777-82.
- 182 Davis JN, et al. Soy isoflavone supplementation in healthy men prevents NF-kappaB activation by TNF-alpha in blood lymphocytes. *Free Radical Biol Med* 2001;30:1293-1302.
- 183 Nagata C, et al. Effect of soymilk consumption on serum estrogen and androgen concentrations in Japanese men. *Cancer Epidemiol Biomarkers Prev* 2001;10:179-84.
- 184 Vacher P, et al. The lipidosterolic extract from *Serenoa repens* interferes with Prolactin receptor signal transduction. *J Biomed Sci* 1995;2:357-65.
- 185 Iguchi k, et al. Myristoleic acid, a cytotoxic component in the extract from *Serenoa repens*, induces apoptosis and necrosis in human prostatic LNCaP cells. *Prostate* 2001;47:59-65.
- 186 Ishii K, et al. Extract from *Serenoa repens* suppresses the invasion activity of human urological cancer cells by inhibiting urokinase-type plasminogen activator. *Biol Pharm Bull* 2001;24:188-90.
- 187 Goldmann WH, et al. Saw palmetto berry extract inhibits cell growth and Cox-2 expression in prostatic cancer cells. *Cell Biol Int* 2001;25:1117-24.
- 188 Debryne F, et al. Comparison of a phytotherapeutic agent (Permixon) with an alpha-blocker (Tamsulosin) in the treatment of benign prostatic hyperplasia: a 1-year randomized international study. *Eur Urol* 2002;41:497-507.
- 189 Mitropoulos D, et al. In vivo effect of the lipidosterolic extract of *Serenoa repens* (Permixon) on mast cell accumulation and glandular epithelium

- trophism in the rat prostate. *World J Urol* 2002;19:457-61.
- 190 Hass, MA, et al. Identification of components of *Prunus Africana* extract that inhibit lipid peroxidation. *Phytomedicine* 1999;6:379-88.
- 191 Wilt T, et al. *Pygeum africanum* for benign prostatic hyperplasia. *Cochrane Database Syst Rev* 2002;1:CD001044.
- 192 Choo MS, et al. Functional evaluation of Tadenan on micturition and experimental prostate growth induced with exogenous dihydrotestosterone. *Urology* 2000;55:292-8.
- 193 Yablonsky F, et al. Antiproliferative effect of *Pygeum africanum* extract on rat prostatic fibroblasts. *J Urol* 1997;157:2381-7.
- 194 Wilt T, et al. Cernilton for benign prostatic hyperplasia. *Cochrane Database Syst Rev* 2000;2:CD001042.
- 195 Roberts KP, et al. Cyclic hydroxamic acid inhibitors of prostate cancer cell growth: selectivity and structure activity relationships. *Prostate* 1998;34:92-9.
- 196 Zhang X, et al. Isolation and characterization of a cyclic hydroxamic acid from a pollen extract, which inhibits cancerous cell growth in vitro. *J Med Chem* 1995;38:735-8.
- 197 Aoki A, et al. [Clinical evaluation of the effect of tamsulosin hydrochloride and cernilton pollen extract on urinary disturbance associated with benign prostatic hyperplasia in a multicentered study]. *Hinyokika Kyo* 2002;48:259-67.
- 198 Kamijo T, et al. Effect of cernilton pollen-extract on experimental nonbacterial prostatitis in rats. *Prostate* 2001;49:122-31.
- 199 Konrad L, et al. Antiproliferative effect on human prostate cancer cells by a stinging nettle root (*Urtica dioica*) extract. *Planta Med* 2000;66:44-7.
- 200 Lamm, DL, Riggs DR. Enhanced immunocompetence by garlic: role in bladder cancer and other malignancies. *J Nutr* 2001;131:1067S-70S.
- 201 Pinto JT, Rivlin RS. Antiproliferative effects of allium derivatives from garlic. *J Nutr* 2001;131:1058S-60S.
- 202 Key TJ, et al. A case-control study of diet and prostate cancer. *Br J Cancer* 1997;76:678-87.
- 203 Hayashi A, et al. Effects of daily oral administration of quercetin chalcone and modified citrus pectin. *Altern Med Rev* 2000;5:546-52.
- 204 Weisburger JH, et al. Mechanisms of chronic disease causation by nutritional factors and tobacco products and their prevention by tea polyphenols. *Food Chem Toxicol* 2002;40:1145-54.
- 205 Satoh K, et al. Inhibition of aromatase activity by green tea extract catechins and their endocrinological effects of oral administration in rats. *Food Chem Toxicol* 2002;40:925-33.
- 206 Bachrach U, Wang YC. Cancer therapy and prevention by green tea: role of ornithine decarboxylase. *Amino Acids* 2002;22:1-13.
- 207 Hiipakka RA, et al. Structure-activity relationships for inhibition of human 5 α -reductases by polyphenols. *Biochem Pharmacol* 2002;63:1165-76.
- 208 Gupta S, et al. Inhibition of prostate carcinogenesis in TRAMP mice by oral infusion of green tea polyphenols. *Proc Natl Acad Sci USA* 2001;98:10350-5.
- 209 Chung Ly, et al. Induction of apoptosis by green tea catechins in human prostate cancer DU145 cells. *Life Sci* 2001;68:1207-14.
- 210 Dorai T, et al. Therapeutic potential of curcumin in human prostate cancer. III. Curcumin inhibits proliferation, induces apoptosis, and inhibits angiogenesis of LNCaP prostate cancer cells in vivo. *Prostate* 2001;47:293-303.
- 211 Ibid.
- 212 Hour TC, et al. Curcumin enhances cytotoxicity of chemotherapeutic agents in prostate cancer cells by inducing p21 (WAF1/CIP1) and C/EBP β expressions and suppressing NF-kappaB activation. *Prostate* 2002;51:211-8.
- 213 Mukhopadhyay A, et al. Curcumin downregulates cell survival mechanisms in human prostate cancer cell line. *Oncogene* 2001;20:7597-7609.
- 214 Rafi MM, et al. Novel polyphenol molecule isolated from licorice root (*Glycyrrhiza glabra*) induces apoptosis, G2/M cell cycle arrest, and Bcl-2 phosphorylation in tumor cell lines. *J Agric Food Chem* 2002;50:677-84.